

Rapid Recovery Hyperbarics

9439 Archibald Ave. Suite 104 Rancho Cucamonga CA, 91730
909.477.4545 | www.hbot4u.com



Birth

Jun 27 2012 3:17 PM

Although stated in technical language, the message is clear: if a child has ANY degree of oxygen deprivation or injury at or before birth, get HBOT as quickly as possible. Further, as this article also points out, early treatment can prevent problems later on that would then be more difficult to reverse. Treat now, and prevent a life time of decreased function.

The Pathogenesis of Cerebral Injury in the Newborn Phillip B James Wolfson
Hyperbaric Medicine Unit The University of Dundee, Scotland, UK.

Birth involves the transfer of the function of oxygenation, the most critical factor in life support from the placental circulation to breathing. This transfer of oxygenation to the lungs of the neonate requires major changes to the neonatal circulation. During labor, placental blood flow is inevitably compromised by uterine contractions, and if oxygenation falls below a critical level, then brain damage may occur at this time. Prystowsky has suggested that mothers should breathe 100% oxygen during labor as a preventative measure, because of the increased gradient for the delivery of oxygen to tissue.

Premature placental separation may also severely compromise fetal oxygenation and delivery may also be associated with fetal brain tissue. In the last few weeks of intrauterine life the cerebral circulation of the fetus matures in preparation for birth. The changes are most prominent in the mid-brain with increased vascularisation and maturation of the blood-brain barrier. If delivery is premature the neonate is at special risk of damage because these critical changes have not yet taken place. Ultrasonic imaging through the anterior fontanelle allowed damage to the midbrain to be detected. Global oxygen deprivation may result in brain atrophy leading to microcephaly, but lesser degrees of hypoxia cause cerebral edema, particularly in the midbrain where the nutrition of some tissue is dependant on the venous drainage. The increased tissue water content reduces oxygen transport and if the edema is not resolved the basal ganglia and the critical areas of the midbrain can undergo cystic

Rapid Recovery Hyperbarics

9439 Archibald Ave. Suite 104 Rancho Cucamonga CA, 91730

909.477.4545 | www.hbot4u.com



degeneration. The blood-brain barrier failure may also be associated with hemorrhage. Damage to the internal capsules in cerebral medulla leads often involves the internal capsules severing the pyramidal tracts. However, in many cases the tracts continue to function for some time but myelination is prevented leaving the axis cylinders exposed. This is associated with a delay in the development of spasticity which may be as long as two years. The additional gradient for the transport of oxygen that is provided by breathing oxygen under Hyperbaric conditions may be able to prevent brain damage in many children as Magnetic Resonance Spectroscopy can detect hypoxia inferred from the presence of lactate. It is possible to screen infants at birth to identify those at risk. Even after a delay, the use of hyperbaric oxygen therapy may be beneficial by resolving the edema. This may allow the completion of myelination and prevent the development of spasticity and global delays or neurological damage.

